

Case 3172***Leptodactylus chaquensis* Cei, 1950 (Amphibia, Anura): proposed conservation of the specific name**

José M. Cei

Departamento Ciencias Naturales, Facultad Ciencias Exactas y Naturales, Universidad Nacional de Río Cuarto, 5800-Río Cuarto, Córdoba, Argentina
(e-mail: fvidela@lab.cricyt.edu.ar)

Abstract. The purpose of this application is to conserve the widely used name *Leptodactylus chaquensis* Cei, 1950 for a subtropical South American frog. The term *typica* was applied in 1948 to what was then supposed to be a 'forma' of *L. ocellatus* Linnaeus, 1758; under Article 45.6.4 of the Code the name *L. typicus* Cei, 1948 is available as a senior synonym of *L. chaquensis*, but it has never been used and its suppression is proposed.

Keywords. Nomenclature; taxonomy; Anura; LEPTODACTYLIDAE; *Leptodactylus*; *Leptodactylus chaquensis*; *Leptodactylus ocellatus*; frogs; South America.

1. In a paper (Cei, 1948) on the seasonal reproductive behaviour of the Argentinian frog *Leptodactylus ocellatus* (Linnaeus, 1758) I described (pp. 308–312) two 'razas' (races) or 'formas', to which I applied, for purposes of discussion, the terms '*typica*' and '*reticulata*'. Other aspects of their reproductive behaviour were described the following year (Cei, 1949).

2. It subsequently became clear that the 'forma' which I had called '*typica*' is a species distinct from *L. ocellatus*, and in a taxonomic account (Cei, 1950) of this group of frogs I established the name *Leptodactylus chaquensis* Cei, 1950 (p. 417) for the taxon. A further discussion of the taxonomy and distribution of the species is given in Cei (1980, pp. 348–352).

3. Nobody has ever used '*typica*' as a valid species-group name, or since Cei (1950) even cited it in the synonymy of *L. chaquensis*. In contrast, the name *L. chaquensis* has been used in a very large number of works, and I give here some examples.

Taxonomic works: Cei, 1980; Frost, 1985; De La Riva, Kohler, Lötters & Reichle, 2000;

Faunistic works: Lynch, 1971; Gallardo, 1979; Gudynas, 1984; Lavilla, Cruz & Scrocchi, 1995; De La Riva & Maldonado, 1999;

Biological and biochemical works: Erspamer, Roseghini & Cei, 1964; Barrio, 1966; Heyer, 1969; Duellman & Trueb, 1986; Roseghini, Erspamer, Falconieri Erspamer & Cei, 1986; Yanosky, Mercolli & Dixon, 1995.

A list of 41 works using the name *Leptodactylus chaquensis* Cei, 1950 has been given to the Commission Secretariat, and more could be supplied.

4. I have become aware, as a result of a message (August 2000) from Dr Darrel Frost, that the terms '*typica*' and '*reticulata*' which I had used in 1948 (see para. 1 above) to refer to two supposed 'formas' (hence their feminine endings) of

Leptodactylus ocellatus are, under Article 45.6.4 of the Code, deemed to be available names established at subspecific rank (contrary to my then intention). The name *typica* (or *typicus*) Cei, 1948 is thus a senior synonym of the specific name of *L. chaquensis* Cei, 1950. However, since *L. chaquensis* has had very extensive usage but nobody has ever used (or even published) the combination *L. typicus* it would be very confusing now to introduce the latter name.

5. The International Commission on Zoological Nomenclature is accordingly asked:

- (1) to use its plenary power to suppress the name *typica* Cei, 1948, as published in the combination *Leptodactylus ocellatus* forma *typica*, for the purposes of the Principle of Priority but not for those of the Principle of Homonymy;
- (2) to place on the Official List of Specific Names in Zoology the name *chaquensis* Cei, 1950, as published in the binomen *Leptodactylus chaquensis*;
- (3) to place on the Official Index of Rejected and Invalid Specific Names in Zoology the name *typica* Cei, 1948, as published in the combination *Leptodactylus ocellatus* forma *typica* and as suppressed in (1) above.

Acknowledgement

I thank Dr Darrel Frost (American Museum of Natural History, New York) for drawing my attention to this matter.

References

- Barrio, A. 1966. Diversidad acustica entre el canto nupcial de *Leptodactylus ocellatus* y *L. chaquensis*. *Physis* (Buenos Aires), **26**: 275–277.
- Cei, J.M. 1948. El ritmo estacional en los fenomenos ciclicos endocrino-sexuales de la rana criolla (*Leptodactylus ocellatus* (L.)). *Acta Zoologica Lilloana*, **6**: 283–331.
- Cei, J.M. 1949. Factores genetico-raciales que diferencian la regulacion hormonal del ciclo sexual en *Lepidodactylus ocellatus* de la Argentina. 'Razas de temperatura' y sus relaciones con algunas características climáticas regionales. *Acta Zoologica Lilloana*, **7**: 113–134.
- Cei, J.M. 1950. *Leptodactylus chaquensis* n. sp. y el valor systematico real de la especie *Leptodactylus ocellatus* en la Argentina. *Acta Zoologica Lilloana*, **9**: 395–423.
- Cei, J.M. 1980. Amphibians of Argentina. *Monografia 2, Monitore Zoologico Italiano*. 599 pp.
- De La Riva, I., Kohler, J., Lötters, S. & Reichle, S. 2000. Ten years of research on Bolivian amphibians: updated checklist, distribution, taxonomic problems, literature and iconography. *Revista Española de Herpetología*, **14**: 19–164.
- De La Riva, I. & Maldonado, M. 1999. First record of *Leptodactylus ocellatus* (Linnaeus, 1758) in Bolivia and comments on related forms. *Graellsia*, **55**: 193–197.
- Duellman, W.E. & Trueb, L. 1986. *Biology of Amphibians*. 669 pp. McGraw Hill, New York.
- Erspamer, V., Roseghini, M. & Cei, J.M. 1964. Indole, imidazole and phenylalkylamines in the skin of thirteen *Leptodactylus* species. *Biochemical Pharmacology*, **13**: 1083–1093.
- Frost, D. 1985. *Amphibian species of the world. Taxonomic and geographical references*. 732 pp. Allen Press, Lawrence, Kansas.
- Gallardo, J.M. 1979. Composición, distribución y origen de la herpetofauna chaqueñas. Pp. 299–307 in Duellman, W.E. (Ed.), *The South American herpetofauna: its origin, evolution and dispersal*. Monograph 7, Museum of Natural History of the University of Kansas.
- Gudynas, E. 1984. Sobre el Rio Uruguay como barrera biogeográfica para Anfibios y la significación de la presencia de *Leptodactylus chaquensis* Cei, 1950 en el Uruguay. *Bolettino della Società Zoologica de Uruguay*, **2**: 78–89.
- Heyer, W.R. 1969. The adaptive ecology of the species groups of the genus *Leptodactylus*. *Evolution*, **23**: 421–428.

- Lavilla, E.O., Cruz, F.B. & Scrocchi, G.J.** 1995. Amphibiens et reptiles de la Station Biologique 'Los Colorados' dans le province de Salta, Argentine. *Revue française d'Aquariologie*, **22**(1-2): 51-128.
- Lynch, J.D.** 1971. Evolutionary relationships, osteology and zoogeography of leptodactylid frogs. *Miscellaneous Publications of the University of Kansas*, no. 53. 238 pp.
- Roseghini, M., Erspamer, V., Falconieri Erspamer, G. & Cei, J.M.** 1986. Indole, imidazole and phenylalkylamines in the skin of one hundred and forty American amphibian species other than Bufonids. *Comparative Biochemistry and Physiology*, **85C**: 139-142.
- Yanosky, A.A., Mercolli, C. & Dixon, J.R.** 1995. Some aspects of the ecology of two sympatric species, *Leptodactylus ocellatus* (Linnaeus, 1758) and *Leptodactylus chaquensis* Cei, 1950, in the humid Chaco of Argentina. *Bulletin of the Maryland Herpetological Society*, **31**(2): 78-92.

Comments on this case are invited for publication (subject to editing) in the *Bulletin*; they should be sent to the Executive Secretary, I.C.Z.N., c/o The Natural History Museum, Cromwell Road, London SW7 5BD, U.K. (e-mail: iczn@nhm.ac.uk).